

# Likelihood II

## Assessment workshop

Arni Magnusson

ICES, 18 Oct 2012

# Outline

- 1 Likelihood - relative probability, support, combine data sources
- 2 Estimation - MLE, log likelihood, confidence interval
- 3 Normal distribution -  $N(\mu, \sigma)$ , dnorm
- 4 Profile likelihood - procedure, interpretation

# Outline

- 1 Likelihood - relative probability, support, combine data sources
- 2 Estimation - MLE, log likelihood, confidence interval
- 3 Normal distribution -  $N(\mu, \sigma)$ , dnorm
- 4 Profile likelihood - procedure, interpretation

# Likelihood concepts

## Relative probability

$$P(A_1) = 0.5 \quad P(A_2) = 0.3 \quad P(A_3) = 0.2$$

$$L(A_1) = 500 \quad L(A_2) = 300 \quad L(A_3) = 200$$

$$L(A_1) = 0.005 \quad L(A_2) = 0.003 \quad L(A_3) = 0.002$$

# Likelihood concepts

Expresses how well the data **support** some parameter value  
or hypothesis

$$L(\theta|\text{data})$$

Like *RSS* but even more useful:  
not only point estimate, but also **uncertainty**

# Likelihood concepts

We can fit a model to many types of data at once and **combine** the likelihood components with simple multiplication

$$L = L_1 \times L_2 \times \dots$$

Unified framework, for simple or complex models

# Likelihood concepts

**Choose** between models with different number of parameters

$$2 \log \frac{L_1}{L_0} \sim \chi^2_{\Delta df}$$


$$\text{AIC} = -2 \log L + 2k$$

$$\text{BIC} = -2 \log L + \log(n)k$$

# Outline

- 1 Likelihood - relative probability, support, combine data sources
- 2 Estimation - MLE, log likelihood, confidence interval
- 3 Normal distribution -  $N(\mu, \sigma)$ , dnorm
- 4 Profile likelihood - procedure, interpretation

# Maximum likelihood estimation



# Log likelihood

**Log** transformation makes things easier

$$L(\theta|\text{data}) = p(\text{data}|\theta)$$




$$p(y_1, \dots, y_n|\theta)$$

$$p(y_1|\theta) \times \dots \times p(y_n|\theta)$$


$$\prod p(y_i|\theta)$$

$$\log L(\theta|\text{data}) = \sum \log p(y_i|\theta)$$


# Log likelihood

**L****log L** **$-\log L$** 


# Confidence interval




# Confidence interval




# Confidence interval




# Confidence interval



# Confidence interval



# Confidence interval



$$0.5\chi_{df=1}^2 = 1.92 \text{ for 95\% confidence interval}$$

# Outline

- 1 Likelihood - relative probability, support, combine data sources
- 2 Estimation - MLE, log likelihood, confidence interval
- 3 Normal distribution -  $N(\mu, \sigma)$ , dnorm
- 4 Profile likelihood - procedure, interpretation

# Normal distribution

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

$$p(y_i|\theta) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(y_i-\mu_i)^2}{2\sigma^2}}$$

$$L(\theta|y) = \prod \left( \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(y_i-\mu_i)^2}{2\sigma^2}} \right)$$

$$-\log L = [0.5n \log(2\pi)] + n \log \sigma + \frac{\sum (y_i - \mu_i)^2}{2\sigma^2}$$


$$= [0.5n \log(2\pi)] + n \log \sigma + \frac{RSS}{2\sigma^2}$$

## dnorm in R


```
L <- prod(dnorm(y, mu, sigma))
```

```
neglogL <- -sum(dnorm(y, mu, sigma, log=TRUE))
```


## dnorm in R



```
dnorm(theta,  
m=150, s=10)
```



```
dnorm(theta,  
m=150, s=10, log=T)
```



```
-dnorm(theta,  
m=150, s=10, log=T)
```

# Outline


- 1 Likelihood - relative probability, support, combine data sources
- 2 Estimation - MLE, log likelihood, confidence interval
- 3 Normal distribution -  $N(\mu, \sigma)$ , dnorm
- 4 Profile likelihood - procedure, interpretation

# Profile likelihood

- 1 Fix  $\theta$  (a parameter of interest) at some value
- 2 Minimize  $-\log L$  by estimating all other parameters
- 3 Save this value of  $-\log L$

Repeat over a range of  $\theta$  values

# Profile likelihood




# Interpretation

Consider a 2-dimensional likelihood surface, describing the likelihood at different values of two parameters:



	$\theta_1 = 1$	$\theta_1 = 2$	$\theta_1 = 3$	$\theta_1 = 4$	$\theta_1 = 5$
$\theta_2 = 1$	5	15	25	20	15
$\theta_2 = 2$	10	40	60	35	20
$\theta_2 = 3$	50	100	75	40	25
$\theta_2 = 4$	10	40	60	35	20
$\theta_2 = 5$	5	15	25	20	15

# Interpretation

Consider a 2-dimensional likelihood surface, describing the likelihood at different values of two parameters:



# Marginal distribution



In likelihood inference, we find the maximum likelihood for each value of  $\theta_1$  across all values of  $\theta_2$

In Bayesian inference, we integrate the likelihood over  $\theta_2$

# Interpretation

Would you place your bet on  $\theta_1=2$  or  $\theta_1=3$  ?

	$\theta_1=1$	$\theta_1=2$	$\theta_1=3$	$\theta_1=4$	$\theta_1=5$	sum
$\theta_2=1$	5	15	25	20	15	80
$\theta_2=2$	10	40	60	35	20	165
$\theta_2=3$	50	100	75	40	25	290
$\theta_2=4$	10	40	60	35	20	165
$\theta_2=5$	5	15	25	20	15	80
sum	80	210	245	150	95	780